


Cancer Survivorship – Tobacco

A survey of 2,938 cancer survivors, conducted and analyzed by American Cancer Society researchers found the below:

- About 1 in 10 survivors still reports smoking about 9 years after a diagnosis
- Survivors who say they smoke consume an average of 15 cigarettes per day—almost a full pack
- Survivors who are young, female, less educated and with lower incomes are the most likely to smoke
- Smoking habits also varied by cancer type, with bladder, lung, and ovarian cancer survivors the most likely to report they currently smoke.
- About 1 in 3 survivors who currently smoke said they intend to quit.


Source: Westmaas, J. L., Alcaraz, K. I., Berg, C. J., & Stein, K. D. (2014). Prevalence and Correlates of Smoking and Cessation-Related Behavior among Survivors of Ten Cancers: Findings from a Nationwide Survey Nine Years after Diagnosis. *Cancer Epidemiology Biomarkers & Prevention*, 23(9), 1783-1792. doi:10.1158/1055-9965.epi-14-0046

Percentage of cancer survivors aged 18 years and older who were current cigarette smokers, Both Sexes, 1992-2017



HP 2020 Target TU-1.1: 12%.
 Source: Centers for Disease Control and Prevention, National Center for Health Statistics. National Health Interview Survey.
 Data are age-adjusted based on the age distribution of cancer patients diagnosed in 2000 in the SEER 18 areas (<http://seer.cancer.gov/registries/terms.html>) using age groups: 18-24, 25-34, 35-44, 45-54, 55-64, 65-74, 75-84, 85+.
 Weighted regression lines are calculated using the Joinpoint Trend Analysis Software, Version 4.6 February 2018, National Cancer Institute.
 The AAPC is the Average Annual Percent Change and is based on the APCs calculated by Joinpoint.
 * The Annual Percent Change (APC)/Average Annual Percent Change (AAPC) is statistically significant.

Trends in secondhand smoke exposure prevalence (%) among cancer survivors in the United States for all nonsmokers and by smoking history



Source: Akinboro, O., Olorunfemi, O., Basak, P., Phillips, E., Pomerantz, D., Bernhardt, B., . . . Ostroff, J. S. (2017). Secondhand Smoke Exposure Among Community-Dwelling Adult Cancer Survivors in the United States: 1999–2012. *Cancer Epidemiology Biomarkers & Prevention*, 26(8), 1296-1305. doi:10.1158/1055-9965.epi-16-0777